

(19) World Intellectual Property Organization
International Bureau




(43) International Publication Date
22 February 2001 (22.02.2001)

PCT

(10) International Publication Number
WO 01/12230 A1

- (51) International Patent Classification⁷: **A61K 47/30**
- (21) International Application Number: **PCT/KR00/00868**
- (22) International Filing Date: **7 August 2000 (07.08.2000)**
- (25) Filing Language: **English**
- (26) Publication Language: **English**
- (30) Priority Data:
1999/33984 17 August 1999 (17.08.1999) **KR**
- (71) Applicants and
(72) Inventors: **PARK, Myung-Ok [KR/KR]; #107-1403**
Hakyeoul Chunggu Apt., Hagye-dong, Nowon-gu, Seoul
139-230 (KR). **LEE, Kang, Choon [KR/KR]; 86-12**
Nonhyeon-dong, Kangnam-ku, Seoul 135-010 (KR).
- (74) Agent: **LEE, Won-Hee; Sung-ji Heights II, 8th floor,**
642-16 Yoksam-dong, Kangnam-ku, Seoul 135-080 (KR).
- (81) Designated States (*national*): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CR, CU, CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW.
- (84) Designated States (*regional*): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).
- Published:
— *With international search report.*
- For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.*

(54) Title: **THE NASAL TRANSMUCOSAL DELIVERY OF PEPTIDES CONJUGATED WITH BIOCOMPATIBLE POLYMERS**



(57) Abstract: The present invention relates to a pharmaceutical composition for a nasal transmucosal delivery, comprising a biocompatible polymer-biologically active peptide conjugate. The pharmaceutical composition of the present invention increases the water solubility of peptide, which is sparingly soluble in water, improves a stability by protecting from being degraded by protease, and, consequently, reduces an administration number of drug to decrease side-effects induced by drug abuse. In addition, since the pharmaceutical composition of the present invention is delivered through the nasal cavity, it allows drug activity to be expressed in a short period of time and improves a bioavailability.

THE NASAL TRANSMUCOSAL DELIVERY OF PEPTIDES
CONJUGATED WITH BIOCOMPATIBLE POLYMERS

FIELD OF THE INVENTION

5

The present invention relates to a pharmaceutical composition for a nasal transmucosal delivery, comprising a biologically active peptide, which is sparingly soluble in water, conjugated with
10 an active biocompatible polymer.

More particularly, the present invention relates to the pharmaceutical composition containing the biocompatible polymer-biologically active peptide conjugate suitable for use in the nasal transmucosal
15 delivery, which is highly improved in water solubility and protected from being degraded by protease.

The pharmaceutical composition comprising of the peptide-polymer conjugate for the nasal transmucosal delivery of the present invention allows drug activity
20 to be expressed in a short period of time and improves a bioavailability.

BACKGROUND

25 In the body, various peptides play important roles, existing as various forms such as hormones and cytokines. With recent great advances in genetic

engineering, various peptides have been able to be synthesized on a mass scale and be used as medicines.

Use of peptides or proteins as medicines, however, suffers from many problems. First, peptides
5 or proteins are very low in body absorption efficiency because they are easily hydrolyzed or degraded by enzymes within a short period of time after being taken into the body. Further, when such peptide medicines are repetitively administered, immune
10 reactions are frequently induced to produce antibodies which may cause so serious hypersensitivity as to menace the life of the administee, acting as a neutralizing role against the physiological activity of the medicines. In addition, the clearance
15 attributable to the reticuloendothelial system (RES) is increased. Therefore, most of peptide medicines have been administered by injection, thus far. Injection administration, however, gives patients pain, accompanying dangers. Particularly, patients who need
20 to be treated for a long period of time may not be able to treat themselves by injection. Thus, there remains a need to develop other routes for peptide administration.

25 An adult's nasal cavity is coated with a mucosa at a thickness of 2.0-4.0 mm (Mugind, Nasal Allergy, Blackwell Scientific, Oxford, 1979) and has a volume

of about 20 ml. It is thought that the nasal cavity allows drug activity to be expressed in a short period of time because of its being abundant in fine villus and large in absorption surface area. Accordingly, extensive research has been made on the transmucosal delivery of drugs. Factors which have influence on the absorption of drugs through mucosae include physical and chemical properties of drugs, such as drug's inherent transmittance, ion strength, flow distribution coefficient and molecular weight, carrier transportation, protease degradation, and physiological conditions of nasal mucosae. In fact, the nasal mucosa is a direct absorption route through which drugs can circumvent the liver metabolism, which is a great hindrance to the utilization of drugs in the body upon oral administration. Thus, the nasal transmucosal route has an advantage over the oral route in that the body utilization of drugs can be significantly improved.

The nasal transmucosal delivery of peptides or proteins of large molecular weights are lower in absorption efficiency than intravenous injection because the peptides or proteins cannot well pass through nasal mucosae. Absorption promoters have been suggested to improve the absorption of peptides. Examples of suggested absorption promoters include surfactants (Hirai et al., *Int. J. Pharm.* 9, 165-169,

1981), acylcarnitine, cholinester, alpha-cyclodextrin, and chelating agents (Lee, In: *Delivery Systems for Peptide Drug*, Plenum, New York, pp 87-104, 1986). These absorption promoters, however, are difficult to
5 apply in practice because they give rise to a decrease in the stability of the drugs upon formulation, or irritate nasal mucosae.

In regard to the nasal transmucosal delivery of
10 drugs, many research results are disclosed in patents.

European Pat. Nos. 23,359 and 122,023 open the possibility that a powder formulation of peptide drugs is delivered through nasal mucosae. U. S. Pat. No. 4,250,163 discloses a mucosa-adsorptive substance
15 which is admixed with a powder form of peptides drugs to enhance the nasal transmucosal delivery of the drugs. European Pat. No. 123,831 is directed to the administration of steroids through nasal mucosae. Germany Pat. No. 2,620,446 describes a body absorption
20 enhancer which is effective for the nasal transmucosal delivery of insulin.

PCT/GB/86/00721 discloses a formulation technique of drugs into microspheres which can be delivered through nasal mucosae. However, this
25 formulation technique can be applied only to particular drugs.

In Japanese Pat. No. Sho. 58-189118,

cyclodextrin is utilized for the nasal transmucosal delivery of peptides. Japanese Pat. No. Sho. 59-89619 discloses ethereal surfactants, for example, polyoxyethylene lauryl ether, as neutral absorption enhancers for nasal transmucosal delivery. However, these surfactants are not suitable for clinical use because they cause damage to nasal mucosae.

Japanese Pat. No. Sho. 61-118325 describes alkaline or neutral amino acids for use in the nasal transmucosal delivery of calcitonin. In Japanese Pat. No. Sho. 63-39822, sucrose fatty acid ester is used as an absorption promoter for the nasal transmucosal delivery of drugs. However, these absorption promoters are also toxic to mucosae.

The above-mentioned reference patents, most of which are based on the sustained release of peptide drugs, enable drugs to be released continuously, but cannot solve the problem in that peptides or protein drugs administered through mucosae are degraded in a short period of time. These conventional techniques find difficulty in being applied for the nasal transmucosal delivery of peptide drugs.

Conjunction of pharmaceutically active proteins or molecules to synthetic macromolecules may afford great advantages when they are applied *in vivo* and *in vitro*. When being covalently bonded to macromolecules,

physiologically active molecules may be changed in surface properties and solubility. For example, an increase may be brought into their solubility in water or organic solvents. Further, the presence of
5 macromolecules may make the conjugated peptides more stable *in vivo* as well as reduce the clearance attributed to the intestinal system, the kidney, the spleen and/or the liver. Hence, conjunction of polymers to peptides can bring about a great
10 improvement in the stability of the peptides in solutions and effectively protect the intrinsic surface properties of peptides to prevent non-specific protein adsorption.

U. S. Pat. No. 4,179,337 discloses conjugates
15 between peptides or polypeptides and polyethylene glycol (hereinafter, referred to as "PEG") with a molecular weight of 500-20,000 or water-soluble polymers, which are reduced in antigenicity and anti-immunity while maintaining the biological activity of
20 the peptides or polypeptides. It is described in U. S. Pat. No. 4,301,144 that hemoglobin is increased in oxygen molecule-carrying potential when being associated with PEG or water-soluble polymers.

Various proteins are reported to show extended
25 half-life spans and reduced immunogenicity in plasma when being conjugated with PEG (Abuchowski et al., *Cancer Biochem. Biophys.*, 7, 175-186, 1984). Uricase-

PEG conjugates are demonstrated to be increased in *in-vivo* half life span and show reduced side-effects during the metabolism of uric acid(Davis et al., *Lancet*, 2, 281-283, 1981).

5 As apparent from the preceding patents, the conjugation of PEG allows biologically active peptides or proteins to be increased in-*vivo* half life span and solubility and to be reduced in immune reactions.

10 Most frequently, the conjugation of PEG to polypeptides is achieved by reacting activated PEG to amino residues of polypeptides. Suitable for use in this purpose are a lysine residue and N-termini. As for PEG activation, one of the hydroxy groups of PEG
15 is substituted with a methyl ether group while the other hydroxy group is bonded to an electrophile functional group (Abuchowski, A. and Davis, F. F. (1981), in *Enzymes as Drugs* (Holsenberg, J. and Roberts, J., eds.)). Examples of activated polymers
20 include PEG-N-hydroxysuccineimide-activated esters, which contain amide bonds, PEG-epoxides and PEG-tresylate, which contain alkyl bonds, PEG-carbonyl imidazole and PEG-nitrophenyl carbonates, which contain urethane bonds, PEG-aldehyde, which contains a
25 Schiff's base at the N-terminus.

On a polypeptide sequence, lysine residues are randomly located, so that PEG is non-specifically

bonded to the polypeptide. In order to obtain uniform PEG-peptide conjugates, there have been made attempts of bonding PEG to targeted sites such as cystein residues, oligo sugars, hydroxy groups, arginine
5 groups.

Examples of PEG derivatives able to specifically react to cystein groups of polypeptides include PEG-vinyl sulphone, PEG-iodoacetamide, PEG-maleimide, and PEG-orthopyridyl disulfide with most preference to
10 maleimide-containing PEG. PEG-vinyl sulfone is best in view of the stability in water solutions while PEG-orthopyridyl disulfide can be reversibly degraded *in vivo* because of the presence of disulfide bonds. Peptides taking advantage of these derivatives can be
15 exemplified by interleukin-3 and interleukin-2.

PEG derivatives reactive specifically to oligo sugars of polypeptides may be exemplified by PEG-hydrizide, which is able to react with aldehyde-containing compounds to form relatively stable
20 hydrozone bonds. Advantage is taken of the specific bonding of PEG-hydrizides to sugar moieties of glycoproteins.

PEG-isocyanates react specifically with hydroxy groups of polypeptides. In order to conjugate PEG to
25 arginine residues of polypeptides, there is used PEG derivatives containing phenylglyoxal, which is highly reactive to the guanidino group.

As mentioned above, the nasal transmucosal delivery of peptides alone is significantly improved in absorption efficiency compared with the oral administration because the peptides are not subjected to liver metabolism, but poor in the bioavailability of the peptides because they are degraded by endogenous enzymes.

To overcome the foregoing and other disadvantages, we, the inventors of the present invention, have developed a pharmaceutical composition for the nasal transmucosal delivery, comprising a sparingly soluble, biologically active polypeptide conjugated with an activated biocompatible polymer. The present invention have confirmed that, when the polymer-peptide conjugate is administered through the nasal cavity, it is improved in water solubility and protected from being degraded by proteases, whereby the medicinal activity of the pharmaceutical composition can be sustained for an extended period of time *in vivo*.

SUMMARY OF THE INVENTION

25

It is an object of the present invention to provide a biologically active peptide-polymer

conjugate suitable for use in a nasal transmucosal delivery.

Further objects and advantages of the present invention will appear hereinafter.

5

The present invention provides a pharmaceutical composition for the nasal transmucosal delivery, comprising a sparingly soluble, biologically active polypeptide conjugated with an activated biocompatible polymer.

10

Further features of the present invention will appear hereinafter.

BRIEF DESCRIPTION OF THE DRAWINGS

15

FIG. 1 represents a result of size exclusion chromatography which isolates sCT and PEG-sCT, where

A ; tri-PEG-sCT,

B ; di-PEG-sCT,

20

C ; mono-PEG-sCT,

D ; sCT.

FIG. 2 shows a pH effect to the production of mono-PEG-sCT that PEG is conjugated with N-terminus of calcitonin, Lys¹⁸ or Lys¹¹, where

25

● ; mono-PEG-sCT(N-terminal conjugate),

□ ; mono-PEG-sCT(Lys¹⁸-conjugate),

△ ; mono-PEG-sCT(Lys¹¹-conjugate).

FIG. 3 shows a molar ratio effect of calcitonin:sCT to the production of mono-PEG-sCT that PEG is conjugated with N-terminus of calcitonin, Lys¹⁸ or Lys¹¹, where

- 5 ● ; mono-PEG-sCT(N-terminal conjugate),
 □ ; mono-PEG-sCT(Lys¹⁸-conjugate),
 △ ; mono-PEG-sCT(Lys¹¹-conjugate).

FIG. 4 shows a reaction time effect of calcitonin:sCT to the production of mono-PEG-sCT that
10 PEG is conjugated with N-terminus of calcitonin, Lys¹⁸ or Lys¹¹, where

- ; mono-PEG-sCT(N-terminal conjugate),
 □ ; mono-PEG-sCT(Lys¹⁸-conjugate),
 △ ; mono-PEG-sCT(Lys¹¹-conjugate).

15 FIG. 5 represents a result of size exclusion chromatography which isolates hPTH and PEG-hPTH(1-34), where

- A ; tri-PEG-hPTH(1-34),
 B ; di-PEG-hPTH(1-34),
20 C ; mono-PEG-hPTH(1-34),
 D ; hPTH(1-34).

FIG. 6 shows a reduction effect of calcitonin concentration in blood when sCT and PEG-sCT conjugate are administered through the nasal cavity, where

- 25 A ; sCT,
 B ; mono-PEG5000-sCTs,
 C ; mono-PEG2000-sCTs,

D ; mono-PEG12000-sCTs.

FIG. 7 shows a calcium reduction effect of isolated mono-PEG-sCT isomers, where

- ; mono-PEG-sCT(N-terminal conjugate, M1),
- 5 ■ ; mono-PEG-sCT(Lys¹⁸-conjugate, M2),
- ◆ ; mono-PEG-sCT(Lys¹¹-conjugate, M3),
- △ ; sCT,
- ; mono-PEG-sCT not isolated its isomers.

10 **DETAILED DESCRIPTION OF PREFERRED EMBODIMENTS**

The term "biocompatible polymers" as used herein means naturally occurring or synthetic compounds which are dissolved in water. By way of example, not
15 limitation, the biocompatible polymers include polyethylene glycol, polypropylene glycol (PPG), polyoxyethylene (POE), polytrimethylene glycol, polylactic acid and its derivatives, polyacrylic acid and their derivatives, polyamino acid, polyvinyl
20 alcohols, polyurethane, polyphosphazene, poly(L-lysine), polyalkylene oxide (PAO), and water-soluble polymers such as polysaccharide, dextran, and non-immunogenic polymers such as polyvinyl alcohol and polyacryl amide.

25 Available in the present invention are the polymers ranging, in molecular weight, from about 200 to 20,000 and preferably from 500 to 12,000.

The present invention provides a peptide-polymer conjugate for a nasal transmucosal delivery, which can be prepared by bonding an activated polymer to a biologically active peptide. In this regard, the bond between the peptides and the polymers may be a covalent bond or a non-covalent bond such as a lipophilic bond or a hydrophobic bond.

In preparing the peptide-polymer conjugates of the present invention, the molar ratio of polypeptides to activated polymers are in the range of about 1:1 to 1:10 and preferably in the range of 1:1 to 1:7. In addition, one to three activated polymers may be conjugated to one polypeptide molecule. Peptide-mono polymer conjugates exert the most effective pharmaceutical activity. In the case of calcitonin-PEG conjugates, PEG can be conjugated to the N-terminus of calcitonin and/or Lys¹⁸ and/or Lys¹¹. Of the resulting calcitonin-PEG conjugates, the calcitonin-(mono)PEG conjugate in which one PEG molecule is conjugated to one calcitonin molecule exhibits the highest calcium reduction effect. A conjugate isomer in which PEG is conjugated to the N-terminus or the Lys¹⁸ shows more sustained and effective calcium reduction activity than does a conjugate isomer in which PEG is conjugated to Lys¹¹. At pH 5, 6 or 7 of the reaction solution, more than

80 % of PEG is conjugated to the N-terminus of calcitonin. On the other hand, at pH 8 or higher, PEG is increasingly conjugated to Lys¹¹ and Lys¹⁸. In contrast, changes in reaction time and molar ratio cannot affect the proportion of isomers.

The binding reaction of the peptide-activated polymer in the present invention is performed in 0.1 M phosphate buffer ranged, in pH, from 6 to 9, at 0 to 25C° of reaction temperature for several minutes to 12 hours.

A method of polymer activation consists of the following steps of; preparing the polymer into polyalkylene oxide(hereinafter, referred to as "PAO") such as monoethoxy-poly(ethylene glycol, mPEG); and changing the other part of PAO into a reaction group having reactivity. The activated polymer forms the peptide-polymer conjugate by reacting with ε-amine group of lysine. Besides the amine group of lysine, carboxy group, activated carbonyl group, oxidized sugar and mercapto group in the peptide can be used as a conjugated moiety.

The present inventors have measured the blood concentration of peptide according to time after the peptide-polymer conjugate is administered to rats through the nasal cavity. Thus, the present inventors have confirmed that the peptide by the nasal transmucosal delivery has a better stability in living

body and sustains its biological activity for a long time.

The peptide of the present invention is not
5 limited to the specific therapeutic agents but applied
to the all substances having biological activity,
particularly, it is desirable to use calcitonin,
parathyroid hormone(hereinafter, referred to as "PTH").
insuline, synthetic enkephalin, growth hormone
10 releasing peptide(hereinafter, referred to as "GHRP"),
leutenizing hormone releasing hormone(hereinafter,
referred to as "LHRH") and its derivatives, secretory
components of hypothalamus, calcitonin gene related
peptide(hereinafter, referred to as "CGRP") and
15 thyroid stimulating hormone and thymic humoral
factor(hereinafter, referred to as "THF").

Calcitonin is a single chain peptide composed of
32 amino acids, forms a ring at N-terminus and has a
20 proline amide group at C-terminus. Calcitonin
inhibits a bone absorption by acting directly to
osteoclast and is used for cure of hypercalcemia,
Pajet's disease, pain from the bone absorption and
osteporosis. Calcitonin is produced in salmon, eel,
25 human, pig, etc., and salmon and eel calcitonin have
the most effect.

Parathyroid hormone (PTH) is a peptide hormone composed of 84 amino acids and secreted from parathyroid. Since parathyroid cell has a recognition site for calcium concentration, PTH secretion
5 increases when calcitonin is lower, and PTH secretion decreases when calcitonin higher. PTH increases a calcium absorption taken from foods at the small intestine, transfers calcium from bone to blood, and ,finally, increases a blood calcium concentration.
10 The major active site of PTH is adrenal cortex. PTH binds to the membrane of adrenal cortex to increase the production of cAMP, IP₃(inositol triphosphate) and DAG(diacyl glycerol).

15 Insulin is a peptide composed of A chain and B chain. A chain composed of 21 amino acids and B chain composed of 30 amino acids are connected by 1 pair of disulfide bond. Disulfide bond exists between 6th amino acid of A chain and 11th amino acid of B chain.
20 When blood sugar level is increased, insulin is immediately excreted from pancreas. Sugar is, then, stored in the form of glycogen by insulin or used as energy source for fat or protein synthesis. In addition, insulin plays an important role in
25 homeostasis of calcium. When insulin presents at a high concentration, calcium uptake is increased from cell exterior to cell interior and hypercalcemia is

induced.

Enkephalin is a pentapeptide representing a similar action to opium. Enkephalin is divided into
5 two groups of methionine enkephalin and leucine enkephalin according to C-terminus bounded to Try-Gly-Gly-Phe. Enkephalin inhibits pain from transferring to brain by preventing a release of substance P from the end of analgesic nerve fiber.

10

Growth hormone releasing peptide(GHRP) presents in the form of hepta and hexa and affects a release of growth hormone. The growth hormone releasing effect of GHRP is related to a dose, increases until
15 adolescence and decreases after then.

GHRP-6 is a hexapeptide having a structure of His-Asp-Try-Ala-Try-Asp-Lys-NH₂, and is very stable in acetate buffer ranged, in pH, from 5.5 to 6.0. In case of oral administration, bioavailability of GHRP-6 is
20 0.3%, an absorption half life is 15 minutes, and an elimination half life is 60 minutes. GHRP-6 secretes selectively growth hormone through a receptor in hypothalamus and pituitary gland.

25 Leuteinizing hormone releasing hormone(LH-RH) is a hypothalamus peptide and stimulates the releases of LH and follicle stimulating hormone. LH-RH regulates a

function of brain and many peripheral organs by binding to receptors on the targeted cell surface. LH-RH has a decapeptide structure of Pyro-Glu-His-Trp-Ser-Tyr-Gly-Leu-Arg-Pro-Gly-NH₂, and is degraded in tubular of the kidney. By way of examples, LH-RH derivatives include nafarelin, busecilin, zilidexin, etc.

Pharmaceutical composition for the nasal transmucosal delivery including the peptide-polymer conjugate of the present invention can be formulated into the suitable form, and mainly administered by spray as a medicine for external use.

For preparation of spray, the peptide-polymer conjugate is dissolved in a solvent, or a suspended, medicinal solution is filled in a container having a specific spraying device(valve) with a low viscous spraying agent. For this, the medicinal solution is sprayed in the type of smog using pressure. The material quality of container uses a metal such as a tinned iron and aluminum.

If necessary, the inside of container is coated with moth-proof painting. When a internal volume is lower than 100 ml, the container made of glass or synthetic resin is possible. The spraying agent uses generally a compressed air and an incombustible liquefied gas, freon(freon 11, CC₁₃F; freon 12, CC₁₂F; freon 114, C₂C₁₂F₄).

A dose of the pharmaceutical compositions containing the peptide-polymer conjugate of the present invention is 0.1 ug - 10 mg/kg/day, and can be widely altered according to the kind of peptide and patient's
5 condition.

EXAMPLES

Practical and presently preferred embodiments of
10 the present invention are illustrative as shown in the following Examples.

However, it will be appreciated that those skilled in the art, on consideration of this disclosure, may make modifications and improvements within the
15 spirit and scope of the present invention.

Example 1: Preparation of PEG-sCT

<1-1> Preparation of PEG5000-sCT

20

According to the method of Abuchowski (Abuchowski et al, *Cancer Biochem. Biophys.*, 7, 175-86, 1984), monomethoxy-poly(ethylene glycol) was prepared from PEG (MW 5000) so that one hydroxyl group of PEG was
25 protected. Phosgene and N-hydroxysuccinimide was added to it to activate in the form of succinyl-N-hydroxysuccinimide ester (hereinafter, referred to as

"SS-PEG"). 4.38 mg of the activated SS-PEG was dissolved in phosphate buffer (pH 8.0), added with 0.2 ml of salmon calcitonin (hereinafter, referred to as "sCT") (Novabiochem, LA Jolla, CA, USA) (5mg/ml, 0.1 M phosphate buffer, pH 8.0), and stirred for 30 minutes at ambient temperature. 0.1 M glycine was added to stop the reaction. Unreacted calcitonin and PEG were removed by analysis using phosphate buffered saline (hereinafter, referred to as "PBS", pH 7.4) to obtain PEG5000-sCT.

<1-2> Preparation of PEG12000-sCT

PEG (MW 12000) was activated with succinimidyl succinate, and PEG12000-sCT was prepared by using 9.2 mg of the activated SS-PEG. All procedures were performed as indicated in the Example <1-1>.

<1-3> Preparation of PEG2000-sCT

PEG (MW 2000) was activated with succinimidyl succinate, and PEG2000-sCT was prepared by using 2 mg of the activated SS-PEG. All procedures were performed as indicated in the Example <1-1>.

Example 2: Isolation of PEG-sCT

PEG-sCT conjugates prepared in the Examples <1-1> to <1-3> were isolated with the aid of size exclusion column (Super HR 12/30, PharmaciaLKB, Sweden) using PBS (pH 7.0) as eluent at 0.4 ml/min of flow rate (FIG. 1).

5 The peptides isolated from the column as a time difference according to the molecular size was isolated into each peak using fluorescence spectrometer (Hitach, Japan). Each isolated fraction was concentrated using Centricon-10 (Amicon, USA) and stored at refrigerator.

10 FIG. 1 represented a fluorescence intensity of each peak measured by fluorescence spectrometer wherein each peak was eluted from the size-exclusion column.

As illustrated in the FIG. 1, tri-PEG-sCT which was conjugated 3 molecules of PEG with 1 molecule of calcitonin and had a biggest molecular size, was
15 firstly eluted from the column and collected within 30 minutes after sample injection into the column. After the tri-PEG-sCT, di-PEG-sCT and mono-PEG-sCT were isolated in succession.

20 In addition, since calsitonin molecule conjugated with PEG had the smallest molecular size, it was eluted last from the column and time for elution took more 40 minutes. That time, the fluorescence intensity of each peak eluted from the column had measured by the
25 fluorescence spectrometer directly connected to the column, and each peak was collected.

Example 3: Preparation of PEG-sCT According to Various pHs, Molar Ratios and Reaction Times

<3-1> Preparation of PEG-sCT According to pH

5

To 88.2 μ l of each of 0.1 M phosphate buffered solutions ranging, in pH, from 5 to 9, 6.8 μ l of calcitonin (10 mg/ml) and 5 μ l of SS-PEG (60 mg/ml) were added and stirred for 30 min at ambient
10 temperature. Addition of 5 μ l of 1 M glycine ceased the reaction between calcitonin and SS-PEG. Unreacted calcitonin and PEG were removed by dialysis using PBS (pH 7.4).

15 **<3-2> Preparation of PEG-sCT According to Molar Ratio of SS-PEG/Calcitonin**

To 6.8 μ l of calcitonin (10 mg/ml), 10 μ l of SS-PEG solutions were added to give mixtures of
20 calcitonin and SS-PEG at molar ratios of 1:1, 1:2, 1:3, 1:5 and 1:10 and these five mixtures were added with 78.2 μ l of a PBS (pH 8.0). From them, PEG-sCT was prepared in the same procedure as in Example <3-1>.

25 **<3-3> Preparation of PEG-sCT According to Reaction Time**

6.8 μ l of calcitonin (10 mg/ml) was added with
10 μ l of an SS-PEG solution to give a mixture of
calcitonin and SS-PEG in a molar ratio of 1:3, and
then, five copies of reaction solutions were prepared
5 by adding to the mixture with 78.2 μ l of PBS (pH 8.0).
Five copies of this solution were allowed to react at
room temperature for 5, 10, 20, 30 and 60 min with
stirring to prepare PEG-sCT as indicated in Example
<3-1>.

10

Example 4: Isolation of Mono-PEG-Calcitonin Isomer

With the aid of size-exclusion columns, PEG-sCT
conjugates prepared in Examples <3-1> to <3-3> were
15 separated into tri-PEG-sCT, di-PEG-sCT and mono-PEG-
sCT. This separation was conducted in a similar
manner to that of Example 2.

The mono-PEG-sCT obtained was divided into three
isomers by reverse phase high performance liquid
20 chromatography (HPLC). In this regard, 100RP-8
(4.0X125 MM, 5 μ M, Merck) used as a column while a
linear gradient of pentafluoropropionic acid (PFPA)-
containing acetonitrile was used as a mobile phase.
The gradient condition was changed with 36-42 %
25 solvent B (0.1 % PFPA added acetonitrile) and 64-58 %
solvent A (0.1% PFPA added distilled water).
Quantitative measurements were made using a UV

absorption meter (215 nm) or a fluorescent absorption meter (excitation 280 nm, emission 315 nm).

Amino acid analysis showed that the three mono-PEG-sCT isomers thus separated were identified to contain PEG conjugated to the N-terminus, Lys¹⁸ and Lys¹¹ of calcitonin, respectively. In addition, it was found that when the reaction solution was 5, 6 or 7 in pH, more than 80 % of PEG was selectively conjugated to the N-terminus. At pH 8 or higher, PEG was increasingly conjugated to the Lys¹¹ and the Lys¹⁸, as shown in Fig. 2. However, there were no differences in the proportion of isomers when the reaction time and the molar ratio were changed, as shown in Figs. 3 and 4.

Example 5: Preparation and Isolation of PEG-PTH(1-34)

<5-1> Preparation and Isolation of PEG5000-PTH(1-34)

50 μ l of SS-PEG5000 (23 mg/0.20 ml, MW 5,000, 0.1 M PBS, pH 7.0) was added with 50 μ l of a PTH solution (2.4 mg/0.15 ml, 0.1 M PBS, pH 7.0) to give a mixture containing a molar ratio of 5:1 of SS-PEG5000:PTH. The mixture was allowed to react for 30 min at ambient temperature with shaking. After 30 min of the reaction, an excess amount of a 0.1 M glycine solution was added to stop the reaction. Unreacted

parathyroid gland hormone and PEG were removed by dialysis using PBS to obtain PEG5000-PTH(1-34).

The PEG 5000-PTH(1-34) conjugate thus prepared was purified by size-exclusion chromatography using an HPLC system and superdex. For this, PBS (pH 7.0) was used as a developing buffer at a flow rate of 0.8 ml/min. Eluted fractions were stored at 4C° until they were measured for UV absorbance. UV absorbance was measured at 215 nm while the excitation and emission waves of fluorescence were fixed in the range of 280 to 315 nm, as shown in Fig. 5.

<5-2> Preparation and Isolation of PEG2000-PTH(1-34)

PEG2000-PTH(1-34) conjugates were prepared and isolated in a similar manner to that of Example <5-1> except that SS-PEG2000 was used at an amount of 2 mg.

Example 6: Preparation and Isolation of PEG-GHRP

20

<6-1> Preparation and Isolation of PEG5000-GHRP-6

50 µl of SS-PEG5000 (23 mg/0.20 ml, MW 5,000, 0.1 M PBS, pH 7.0) was added with 50 µl of a GHRP-6 solution (0.6 mg/0.15 ml, 0.1 M PBS, pH 7.0) to give a mixture containing a molar ratio of 5:1 of SS-PEG: GHRP-6, prepared PEG-GHRP-6, and isolated and purified

as indicated in the Example <5-1>. Isolated mono-PEG-GHRP-6 was immediately frozen-dried by freeze dryer.

<6-2> Preparation and Isolation of PEG2000-GHRP-6

5

Except for use of PEG2000, PEG2000:GHRP-6 conjugate as indicated in the Example <6-1>.

Example 7: Preparation and Isolation of PEG-LHRH

10

<7-1> Preparation and Isolation of PEG5000-LHRH

50 μ l of SS-PEG5000 (23 mg/0.20 ml, MW 5,000, 0.1 M PBS, pH 7.0) was added with 50 μ l of a LHRH solution (0.5 mg/0.1 ml, 0.1 M PBS, pH 7.0) to give a mixture containing a molar ratio of 1:5 of LHRH:SS-PEG, prepared PEG-LHRH, and isolated and purified as indicated in the Example <5-1>.

20

<7-2> Preparation and Isolation of PEG2000-LHRH

Except for use of PEG2000, PEG2000-LHRH conjugate was prepared, isolated and purified as indicated in the Example <7-1>.

25

Example 8: Preparation and Isolation of PEG5000-triptorelin

Except for use of triptorelin, LHRH derivative, instead of LHRH, PEG5000-triptorelin conjugate was prepared, isolated and purified as indicated in the
5 Example <7-1>.

Example 9: Preparation and Isolation of PEG5000-orntide

10 Except for use of orntide, LHRH agonist, instead of LHRH, PEG5000-orntide conjugate was prepared, isolated and purified as indicated in the Example <7-1>.

Experimental Example 1: Comparison of Calcium Reduction
15 **Effect of sCT and PEG-sCT**

SD (Sprague Dawley) male rat (220 - 300 g, Charles River Japan, Atsugi, Japan) was put under anesthesia by administrating 45 mg/kg of peptobarbital
20 into interperitoneal. Bronchus and esophagus were cannulated with PE-250 tube and thigh arteria with SP-45 tube. 20 ul of a placebo and experimental drugs were administered into the nasal cavity of rats with Hamilton syringe. The placebo used physiological
25 saline and the experimental drug used PEG5000-sCT, PEG12000-sCT, PEG2000-sCT and sCT prepared from the Example 2. Salmon calcitonin was administered at dose

of 0.05 to 4.5 IU per rat, and rats were grouped into 4. 200 ul of blood was taken at before, 5, 10, 30, 60, 120, 240, 480 and 360 minutes after the placebo and the experimental drugs were administered, and immediately
5 centrifuged to obtain 100 ul of plasma. The calcium concentration in plasma was measured using a calcium assay kit (Sigma, USA), and calcium remaining efficiency was calculated as the calcium concentration before administration was 100. The unit of the
10 measured calcium concentration in plasma was mg/dl, and the remaining efficiency was calculated as the ratio (%) to the primary value.

After the remaining coefficient was calculated according to each time, area-under the curve (AUC) was
15 calculated about the average value of remaining coefficient at each time per experimental group. The reduction ratio of calcium concentration was calculated by the mathematical formula 1, and the results were represented in the table 1 (FIG. 6).

20

<Mathematical formula 1>

$$\Delta D (\%) = (AUC_{Ca-plac} - AUC_{Ca-sCT}) / AUC_{Ca-plac} \times 100$$

25 where, $AUC_{Ca-plac}$ was AUC from placebo administration to 960 minutes, and AUC_{Ca-sCT} was AUC from calcitonin and experimental drugs administration to 960

minutes.

<Table 1> The reduction ratio of calcitonin concentration in plasma at each time (n=4)

5

Time	sCT	PEG5000-sCT	PEG12000-sCT	PEG2000-sCT
0	100%	100%	100%	100%
5	98.9%	100.3%	96.8%	97.9%
10	93.7%	97.7%	96.2%	94.5%
30	95%	93.6%	95.0%	91.5%
60	98.7%	86.9%	97.0%	88.0%
120	97.8%	82.2%	98.4%	84.8%
240	100.3%	73.9%	101.3%	82.2%
480	101.6%	82.3%	-	73.9%
960	101%	85.8%	101.8%	71.2%

As illustrated in the table 1, the calcium concentration of calcitonin conjugated without any PEG was decreased for 1 hour after the nasal transmucosal administration, but restored at the original concentration after 2 hours. However, when calcitonin conjugated with PEG was administered, the calcitonin reduction effect represented for 4 hours to 8 hours after the nasal transmucosal administration. In addition, the calcium reduction effect was significantly altered according to the molecular size

of PEG.

Namely, in case of lower 5,000 of molecular weight, the calcium reduction effect was similar sustained, but, in case of 12,000, the calcium reduction effect was not sufficient. This result was because that the more the molecular weight of PEG was increased, the more the degree of absorption into body through the nasal transmucosal was decreased. Consequently, calcitonin conjugated with the lower molecular weight of PEG had a better calcitonin reduction effect than calcitonin conjugated without any PEG, and its calcium reduction effect was sustained for a long time.

Therefore, the nasal transmucosal delivery using the PEG conjugates of the present invention could reduce a used amount of calcitonin and side effects of drugs.

Experimental Example 2: Comparison of Calcium Reduction Effect of Mono-PEG-sCT Isomers

The calcium reduction effect of the mono-PEG-sCT isomers prepared from the Example 4 was measured as indicated in the Example 1 (FIG. 7).

As illustrated in the FIG. 7, isomers conjugated with PEG at N-terminus and Lys¹⁸ represented a better sustaining and effective calcium reduction effect than

isomers conjugated with PEG at Lys¹¹. In addition, in case of mono-PEG-sCT not isolated, Lys¹⁸ isomers having a lower calcium reduction effect than N-terminus isomers represented a considerable reduction effect to sCT or Lys¹¹ isomers.

Example 10: Preparation and Isolation of PEG2000-GLP-1(glucagons like peptide)

PEG (MW 2000) was activated with succinimidyl succinate, and PEG2000-GLP-1 was prepared by using 2 mg of the activated SS-PEG. All procedures were performed as indicated in the Example <1-1>. In addition, PEG2000-GLP-1 was isolated and purified as indicated in the Example <5-1>.

INDUSTRIAL APPLICABILITY

A pharmaceutical composition comprising a peptide-polymer conjugate for a nasal transmucosal delivery of the present invention increases the water solubility of peptide, which is sparingly soluble in water, improves a stability by protecting from being degraded by protease, and, consequently, reduces an administration number of drug to decrease side-effects induced by drug abuse.

In addition, since the pharmaceutical composition

comprising a peptide-polymer conjugate for the nasal transmucosal delivery of the present invention is delivered through the nasal cavity, it allows drug activity to be expressed in a short period of time and
5 improves a bioavailability.

Those skilled in the art will appreciate that the conceptions and specific embodiments disclosed in the foregoing description may be readily utilized as a
10 basis for modifying or designing other embodiments for carrying out the same purposes of the present invention. Those skilled in the art will also appreciate that such equivalent embodiments do not depart from the spirit and scope of the invention as set forth in the appended
15 claims.

What is Claimed is

1. A pharmaceutical composition for a nasal transmucosal delivery which is conjugated a
5 biologically active peptide with an active biocompatible polymer.
2. The pharmaceutical composition for the nasal transmucosal delivery according to claim 1,
10 wherein the peptide is selected from the group of calcitonin, parathyroid hormone (PTH), insulin, synthetic enkephalin, growth hormone releasing peptide (GHRP), luteinizing hormone releasing hormone (LHRH) and its derivatives, secretory
15 components of hypothalamus, calcitonin gene related peptide (CGRP), thyroid stimulating hormone and thymic humoral factor (THF).
3. The pharmaceutical composition for the nasal
20 transmucosal delivery according to claim 1, wherein the biocompatible polymer has 200-20,000 of the molecular weight.
4. The pharmaceutical composition for the nasal
25 transmucosal delivery according to claim 1, wherein the biocompatible polymer has 500-12,000 of the molecular weight.

5. The pharmaceutical composition for the nasal transmucosal delivery according to claim 1, wherein the biocompatible polymer is selected one from the group of polyethylene glycol, polypropylene glycol (PPG), polyoxyethylene (POE), polytrimethylene glycol, polylactic acid and its derivatives, polyacrylic acid and their derivatives, polyamino acid, polyvinyl alcohols, polyurethane, polyphosphazene, poly(L-lysine), polyalkylene oxide (PAO), and water-soluble polymers such as polysaccharide, dextran, and non-immunogenic polymers such as polyvinyl alcohol and polyacryl amide.

15

6. The pharmaceutical composition for the nasal transmucosal delivery according to claim 1, wherein a molar ratio of the peptide and the active biocompatible polymer is 1:1 to 1:7.

20

7. The pharmaceutical composition for the nasal transmucosal delivery according to claim 1, wherein a reaction pH of the peptide and the active biocompatible polymer is 5.0 to 8.0.

25

8. The pharmaceutical composition for the nasal transmucosal delivery according to claim 1, which

is conjugated 1 to 3 molecules of the active biocompatible polymer with 1 molecule of the peptide.

5 9. The pharmaceutical composition for the nasal transmucosal delivery according to claim 8, which is conjugated 1 to 3 molecules of PEG with 1 molecule of calcitonin.

10 10. The pharmaceutical composition for the nasal transmucosal delivery according to claim 9, wherein PEG conjugated at N-terminus, Lys¹⁸ or Lys¹¹ of calcitonin.

15

FIG. 1




FIG. 2




FIG. 3




FIG. 4




FIG. 5




FIG. 6





FIG. 7



INTERNATIONAL SEARCH REPORT

International application No.

PCT/KR00/00868

A. CLASSIFICATION OF SUBJECT MATTER**IPC7 A61K 47/30**

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

A61K 47/30

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

CA Online

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	WO 9852614 (THE BOARD OF TRUSTEES OF THE LELAND STANFORD JUNIOR UNIVERSITY) 26. November 1998 (26. 11. 98)	1-10
A	US 5359030 (Protein Delivery Inc.) 25. October 1994 (25. 10. 94)	1-10



Further documents are listed in the continuation of Box C.



See patent family annex.

* Special categories of cited documents:

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier application or patent but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art

"&" document member of the same patent family

Date of the actual completion of the international search

29 SEPTEMBER 2000 (29.09.2000)

Date of mailing of the international search report

30 SEPTEMBER 2000 (30.09.2000)

Name and mailing address of the ISA/KR

Korean Industrial Property Office
Government Complex-Taejon, Dunsan-dong, So-ku, Taejon
Metropolitan City 302-701, Republic of Korea

Facsimile No. 82-42-472-7140

Authorized officer

YOON, Kyoung Aei

Telephone No. 82-42-481-5609



INTERNATIONAL SEARCH REPORT

Information on patent family members

International application No.

PCT/KR00/00868

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 9852614 A2	26. 11. 98	AU 9875938 A1 EP 975370 A2	11. 12. 98 02. 02. 00
US 5359030	25. 10. 94	WO 9426778 A1 CA 2162366 AA EP 707596 A1 JP 08510255 T2 US 5438040 A US 5681811 A	24. 11. 94 24. 11. 94 24. 04. 96 29. 10. 96 01. 08. 95 28. 10. 97